

**LIST OF APPROVED DRUG FROM 01-01-2012 to Till Date**

<b>S.No</b>	<b>Name Of Drug</b>	<b>Indication</b>	<b>Date of issue</b>
1.	Ornidazole IP 1 % w/w + Povidone Iodine IP (0.5% w/w available Iodine) 5% w/w Ointment	For the prevention and treatment of surgical wound infections in adult patients undergoing surgery	03.01.12
2.	Aceclofenac IP (Sustained released pellets) 200mg/200mg + Thiocolchicoside IP (Sustained released pellets) 8mg/16mg Capsules. (New Dosage Form)	For the treatment of acute inflammation conditions associated with spasm in adults only	04.01.12
3.	Budesonide IP 1mg/0.5mg + Formoterol Fumarate Dihydrate IP eq to Formoterol Fumarate 20mcg/20mcg Respirator suspension. (Additional Indication)	In the regular treatment of patients with moderate to severe Chronic Obstructive Pulmonary Disease (COPD) with frequent symptoms and a history of exacerbations.	04.01.12
4.	Diclofenac Sodium IP (Enteric coated granules) 50mg + Eperisone Hcl IP (Tablet) 50mg Hard Gelatin Capsules. (New Dosage Form)	For the treatment of Patients with acute musculoskeletal spasm associated with low back pain	06.01.12
5	Degarelix for Injection 80mg / 120mg per vial	For the treatment of adult patients with advanced hormone dependent prostate cancer.	19.01.12
6	Linagliptin film coated Tablet 5mg	As an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.  Important limitations of use: 1. Should not be used in patients with type 1 diabetes or for the treatment of diabetic ketoacidosis. 2. Has not been studied in combination with insulin.	23.01.12

7	Eribulin mesylate solution for injection 0.88mg / 2ml vial	For the treatment of patients with locally advanced or metastatic breast cancer who have progressed after at least two chemotherapeutic regimens for advanced disease. Prior therapy should have included an anthracycline and a taxane unless patients were not suitable for these treatments.	27.01.12
8	Plerixafor Injection 20mg/ml	In combination with Granulocyte Colony Stimulating Factor (G-CSF) to enhance mobilisation of hematopoietic stem cells to the peripheral blood for collection and subsequent autologous hematopoietic stem cells transplantation in patients with non-Hodgkin's lymphoma and multiple myeloma whose cells mobilise poorly.	31.01.12
9	Cephalexin IP (as monohydrate) 200mg + Kanamycin IP (as monosulphate) 100,000 IU Suspension per 10 gm Intramammary Suspension	For the treatment of clinical mastitis in lactating dairy cows for bacteria susceptible to the combination of cephalexin and kanamycin such as Staphylococcus aureus, Streptococcus dysgalactiae, Streptococcus uberis and Escheria coli.	07.03.12